

(G, m) -Multiparking Functions

Hungyung Chang^a Po-Yi Huang^{b,*} Jun Ma^{c,†} Yeong-Nan Yeh^{d,‡}

^{a,c,d} Institute of Mathematics, Academia Sinica, Taipei, Taiwan

^b Department of Mathematics, National Cheng Kung University, Tainan, Taiwan

Abstract

The conceptions of G -parking functions and G -multiparking functions were introduced in [15] and [12] respectively. In this paper, let G be a connected graph with vertex set $\{1, 2, \dots, n\}$ and $m \in V(G)$. We give the definition of (G, m) -multiparking function. This definition unifies the conceptions of G -parking function and G -multiparking function. We construct bijections between the set of (G, m) -multiparking functions and the set of $\mathcal{F}_{G,m}$ of spanning color m -forests of G . Furthermore we define the (G, m) -multiparking complement function, give the reciprocity theorem for (G, m) -multiparking function and extend the results [25, 12] to (G, m) -multiparking function. Finally, we use a combinatorial methods to give a recursion of the generating function of the sum $\sum_{i=1}^n a_i$ of G -parking functions $(a_1, \dots, a_n)$.

Keywords: parking functions; spanning forest

1 Introduction

J. Riordan [17] define the parking function as follows: m parking spaces are arranged in a line, numbered 1 to n left to right; n cars, arriving successively, have initial parking preferences, a_i for i , chosen independently and at random; $(a_1, \dots, a_n)$ is called preference function; if space a_i is occupied, car i moves to the first unoccupied space to the right; if all the cars can be parked, then the preference function is called parking function.

Konheim and Weiss [11] introduced the conception of the parking functions of length n in the study of the linear probes of random hashing function. J. Riordan [17] studied the parking functions and derived that the number of parking functions of length n is $(n+1)^{n-1}$, which coincides with the number of labeled trees on $n+1$ vertices by Cayley's formula. Several bijections between the two sets are known (e.g., see [7, 17, 18]). Parking functions have been found in connection to many other combinatorial structures such as acyclic mappings, polytopes, non-crossing partitions, non-nesting partitions, hyperplane arrangements, etc. Refer to [8, 7, 9, 16, 19, 20] for more information.

*Partially supported by NSC 96-2115-M-006-012

†Email address of the corresponding author: majun@math.sinica.edu.tw

‡Partially supported by NSC 96-2115-M-001-005

Parking function $(a_1, \dots, a_n)$ can be redefined that its increasing rearrangement $(b_1, \dots, b_n)$ satisfies $b_i \leq i$. Pitman and Stanley generalized the notion of parking functions in [16]. Let $\mathbf{x} = (x_1, \dots, x_n)$ be a sequence of positive integers. The sequence $\alpha = (a_1, \dots, a_n)$ is called an $\mathbf{x}$ -parking function if the non-decreasing rearrangement $(b_1, \dots, b_n)$ of α satisfies $b_i \leq x_1 + \dots + x_i$ for any $1 \leq i \leq n$. Thus, the ordinary parking function is the case $\mathbf{x} = (1, \dots, 1)$. By the determinant formula of Gončarov polynomials, Kung and Yan [14] obtained the number of $\mathbf{x}$ -parking functions for an arbitrary $\mathbf{x}$. See also [23, 24, 25] for the explicit formulas and properties for some specified cases of $\mathbf{x}$.

Recently, Postnikov and Shapiro [15] gave a new generalization, building on work of Cori, Rossin and Salvy [1], the G -parking functions of a graph. For the complete graph $G = K_{n+1}$, the defined functions in [15] are exactly the classical parking functions. Chebikin and Pylyavskyy [2] established a family of bijections from the set of G -parking functions to the spanning trees of that graph. Then Dimitrije Kostic and Catherine H. Yan [12] proposed the notion of a G -multiparking function, a natural extension of the notion of a G -parking function and extended the result of [25] to arbitrary graphs. They constructed a family of bijections from the set of G -multiparking functions to the spanning forests of G . By the definition in [12], it is easy to see that the vertex 1 is always the root in all G -multiparking functions f if the vertex set of G is $\{1, 2, \dots, n\}$. One of the motivations of this paper is to consider the case in which the vertex 1 isn't the root. So, we give the definition of (G, m) -multiparking function. This definition unifies the conceptions of G -parking function and G -multiparking function because a (G, m) -multiparking function is a G -multiparking functions and a G -parking functions when $m = 1$ and $m = n$ respectively. Using the methods developed by Dimitrije Kostic and Catherine H. Yan [12], we construct bijections between the set $\mathcal{MP}_{G,m}$ of (G, m) -multiparking functions and the set of $\mathcal{F}_{G,m}$ of spanning color m -forests of G .

Richard Stanley's book [22], in the context of rational generating functions, devotes an entire section to exploring the relationships (called reciprocity relationships) between positively- and nonpositively-indexed terms of a sequence. If $\alpha = (a_1, a_2, \dots, a_n)$ is a K_{n+1} -parking function of length n , then $(n - a_1, n - a_2, \dots, n - a_n)$ is called a complement of the parking function α . It is easy to see that the sums $\sum_{i=1}^n (n - a_i) - \binom{n+1}{2}$ and $\sum_{i=1}^n a_i$ are connected with the reciprocity law for K_{n+1} -parking function. Also the sum $\sum_{i=1}^n (n - a_i) - \binom{n+1}{2}$ is one of the most important statistic of K_{n+1} -parking function. Knuth [10] indicated that it corresponds to the number of linear probes in hashing functions. Kreweras [13] concluded that it is equal to the number of inversions in labeled trees on $[n + 1]$. Also it coincides with the number of hyperplanes separating a given region from the base region in the extended Shi arrangements [21]. Catherine H. Yan [25] gave a combinatorial explanation, which revealed the underlying correspondence between the classical parking functions and labeled, connected graphs. Furthermore, for arbitrary graph G , Dimitrije Kostic and Catherine H. Yan [12] indicate the relations between G -inversions and G -multiparking functions. There is a interesting problem: how to define the complement of a (G, m) -multiparking function? In this

paper, we define the (G, m) -multiparking complement function, give the reciprocity theorem for (G, m) -multiparking function and extend the results [25, 12] to (G, m) -multiparking function.

In [12], Dimitrije Kostic and Catherine H. Yan related G -multiparking functions to the Tutte polynomial $T_G(x, y)$ of G . The generating function $P_G(q)$ of G -parking functions is defined as $P_G(q) = \sum_f q^{\sum_{i \in V(G)} f(i)}$ where f ranges over all G -parking functions. Their results implies that $P_G(q) = q^{|E(G)| - |V(G)|} T_G(1, \frac{1}{q})$. Note that the Tutte polynomial of G satisfies the recursion

$$T_G(x, y) = \begin{cases} xT_{G-e}(x, y) & \text{if } e \text{ is a bridge} \\ yT_{G-e}(x, y) & \text{if } e \text{ is a loop} \\ T_{G-e}(x, y) + T_{G \setminus e}(x, y) & \text{otherwise} \end{cases}$$

where $G - e$ is a graph obtained by deleting the edge e and $G \setminus e$ is a graph obtained from G contracting the the vertices i and j from G . By a combinatorial method, we give a recursion of the generating function of the sum $\sum_{i=1}^n a_i$ of G -parking functions $(a_1, \dots, a_n)$.

This paper is organized as follows. In Section 2, we construct bijections between the set $\mathcal{MP}_{G,m}$ of (G, m) -multiparking functions and the set of $\mathcal{F}_{G,m}$ of spanning color m -forests of G . In Section 3, we define the (G, m) -multiparking complement function and study its properties. In Section 4, by a combinatorial method, we obtain a recursion of the generating function of the sum $\sum_{i=1}^n a_i$ of G -parking functions $(a_1, \dots, a_n)$.

2 (G, m) -multiparking functions

In this section, first we give the definition of (G, m) -multiparking function. Given a $m \in \{1, 2, \dots, n\}$, for any $I \subseteq [n]$, let $\alpha(I, m) = \min\{i \in I \mid i \geq m\}$.

Definition 2.1 Let $1 \leq m \leq n$ and G be a connected graph with vertex set $V(G) = \{1, 2, \dots, n\}$. A (G, m) -multiparking function is a function $f : V(G) \rightarrow \mathbb{N} \cup \{-1\}$, such that for every $I \subseteq V(G)$ either (A) $f(\alpha(I, m)) = -1$, or (B) there exists a vertex $i \in I$ such that $0 \leq f(i) < \text{outdeg}_I(i)$.

The vertices which satisfy $f(i) = -1$ in (A) will be called roots of f . Furthermore, we say that the vertex v is called a *absolute root* if $f(v) = -1$ in all (G, m) -multiparking functions f ; and the vertex v is called a *relative root* if there are (G, m) -multiparking functions f and f' such that $f(v) = -1$ and $f'(v) \geq 0$ respectively. By Definition 2.1, it is easy to see that a (G, m) -multiparking function is a G -multiparking functions and a G -parking functions when $m = 1$ and $m = n$ respectively.

Let G be a connected graph with vertex set $V(G) = [n]$, where $[n]$ denote the set $\{1, 2, \dots, n\}$. There are loops and multiple edges in G . Given $m \in [n]$, let m be the absolute root of G . For any $i, j \in [n]$, let $\mu_G(i, j)$ be the number of edges between the vertices i and j in G . For establishing the bijections, all edges of G are colored. The colors of edges connecting the vertices i and j are $0, 1, \dots, \mu_G(i, j) - 1$ respectively for any $i, j \in V(G)$. We use $\{i, j\}_k$ to denote the edge $e \in E(G)$

connecting two vertices i and j with color k . A color m -subforest F of G is a color subgraph of G without cycles such that there is a vertex $j \in [m, n]$ in every component of F . Let $\mathcal{MP}_{G,m}$ and $\mathcal{F}_{G,m}$ be the sets of the (G, m) -multiparking functions and the spanning color m -forests of G respectively. For any $F \in \mathcal{F}_{G,m}$ and $e \in F$, let $c_F(e)$ denote the color of edge e in F . By modifying the algorithms A and B in [12], we construct the bijection Φ between $\mathcal{MP}_{G,m}$ and $\mathcal{F}_{G,m}$. Since the proof of the bijection is similar to the proof in [12], we only give the sketch of the proof in this paper. The following algorithm gives a mapping Φ from $\mathcal{MP}_{G,m}$ to $\mathcal{F}_{G,m}$.

Algorithm A. (Kostic, Yan [12])

Step 1: Let $val_0 = f$, $P_0 = \emptyset$, $F_0 = Q_0 = \{m\}$.

Step 2: At time $i \geq 1$, let $v = \min\{\tau(w) \mid w \in Q_{i-1}\}$, where τ is a vertex ranking in S_n .

Step 3: Let $N = \{w \notin P_{i-1} \mid 0 \leq val_i(w) \leq \mu(w, v) - 1 \text{ and } \{w, v\}_{val_i(w)} \in E(G)\}$ and $\hat{N} = \{w \notin P_{i-1} \mid val_i(w) \geq \mu(w, v) \text{ and } \{w, v\}_{val_i(w)} \in E(G)\}$. Set $val_i(w) = val_{i-1}(w) - \mu_G(w, v)$ for all $w \in \hat{N}$. For any other vertex u , set $val_i(w) = val_{i-1}(w)$. Update P_i , Q_i and F_i by letting $P_i = P_{i-1} \cup \{v\}$, $Q_i = Q_{i-1} \cup N \setminus \{v\}$ if $Q_{i-1} \cup N \setminus \{v\} \neq \emptyset$, otherwise $Q_i = \{u\}$ where $u = \alpha([n] \setminus P_i, m)$. Let F_i be a graph on $P_i \cup Q_i$ whose edges are obtained from those of F_{i-1} by joining edges $\{w, v\}_{val_{i-1}(w)}$ for each $w \in N$.

Define $\Phi = \Phi_{G,m,\tau} : \mathcal{MP}_{G,m} \rightarrow \mathcal{F}_{G,m}$ by letting $\Phi(f) = F_n$. First, iterating the Steps 2-3 until n , we must have $P_n = [n]$ and $Q_n = \emptyset$. Otherwise, let $\tilde{P} = [n] \setminus P_n$. Then $f(i) \geq outdeg_{\tilde{P}}(i)$ for all $i \in \tilde{P}$, a contradiction. Also, it is easy to see that each F_i is a forest since every edge $\{v, w\}_k$ in $F_i \setminus F_{i-1}$ has one endpoint in $V(F_i) \setminus V(F_{i-1})$. In the above algorithm, let f be a (G, m) -multiparking function. Then each tree component T of $\Phi(f)$ has exactly one vertex v with $f(v) = -1$. In particular, $v = \alpha(V(T), m)$. Thus, we have $\Phi(\mathcal{MP}_{G,m}) \subseteq \mathcal{F}_{G,m}$.

Let $F \in \mathcal{F}_{G,m}$. Suppose T is a component of F . Let $\alpha(V(T), m)$ be the root of T . For any non-root $v \in [n]$, there is a unique root r_i which is connected with v . Define the height of v to be the number of edges in the path connecting v with root r_i . If the height of a vertex w is less than the height of v and $\{v, w\}_k$ is an edge of F , then w is the predecessor of v , v is a child of w , and write $w = pre_F(v)$ and $v \in child_F(w)$. The following algorithm will give the inverse map of Φ .

Let G be a connected color graph with a spanning color m -subforest F . A leaf of F is a vertex $v \in V(F)$ with degree 1 in F . Denote the set of leaves of F by $Leaf(F)$. Let $T_1, T_2, \dots, T_k$ be the trees of F with respective roots $m = r_1 < r_2 < \dots < r_k$, where $r_i = \alpha(V(T_i), m)$.

Algorithm B (Kostic, Yan [12]).

Step 1. Let τ be a vertex ranking in S_n . Assume $v_1, v_2, \dots, v_i$ are determined, where $v_1 = m$. Let $V_i = \{v_1, v_2, \dots, v_i\}$ and $W_i = \{v \notin V_i \mid \{v, w\} \in F \text{ for some } w \in V_i\}$.

Case (1) If $W_i = \emptyset$, let v_{i+1} be the minimum vertex which is larger than or equal to m in the set $V(G) \setminus V_i$;

Case (2) Otherwise, let F' be the forest obtained by restricting F to $V_i \cup W$. Let $v_{i+1} = \min\{\tau(w) \mid w \in Leaf(F')\}$.

Step 2. Use $\sigma(F)$ to denote the number of the connected components in F . Set $f(r_1) = f(r_2) = \dots = f(r_{\sigma(F)}) = -1$. For any other vertex v , let $f(v)$ be equal to the sum of the color of the edge connecting the vertices v with $pre_F(v)$ and the cardinality of the set $N(v)$, where $N(v) = N_{G,F,\tau}(v) = \{v_j \mid \{v, v_j\}_k \in E(G) \text{ and } \pi^{-1}(v_j) < \pi^{-1}(pre_F(v))\}$.

Define $\Psi = \Psi_{G,m,\tau} : \mathcal{F}_{G,m} \rightarrow \mathcal{MP}_{G,m}$ by letting $\Psi_{G,m,\tau}(f) = f_F$. For any $I \subseteq [n]$, Let π be the permutation defined in Step 1 of the Algorithm B. It is easy to see that there exists a unique integer k such that $I \subseteq \{\pi(k), \dots, \pi(n)\} = I'$ and $\pi(k) \in I$. If the vertex $\pi(k)$ isn't the root of a component in F , then we have $0 \leq f(\pi(k)) < outdeg_{I'}(\pi(k))$. So, $f(\pi(k)) < outdeg_I(\pi(k))$ since $outdeg_I(\pi(k)) \leq outdeg_{I'}(\pi(k))$. If the vertex $\pi(k)$ is the root of a component in F , then $f(\pi(k)) = -1$ and $\pi(k) = \alpha(I', m)$. Clearly, $\pi(k) = \alpha(I, m)$ since $I \subseteq I'$. Hence, by the Definition 2.1, we have $f = f_F$ is a (G, m) -multiparking function and $\Psi(\mathcal{F}_{G,m}) \subseteq \mathcal{MP}_{G,m}$.

Finally, note that the order $\pi = v_1 v_2 \dots v_n$ in the algorithm B is exactly the order in which vertices of G will be placed into the set P_i when running algorithm A on f . Thus, we have $\Psi(\Phi(f)) = f$. Hence, $\Phi = \Phi_{G,m,\tau}$ and $\Psi = \Psi_{G,m,\tau}$ are inverses of each other. We state these results as the following theorem.

Theorem 2.1 (*Kostic, Yan [12]*) *The mapping Φ is a bijection from $\mathcal{MP}_{G,m}$ to $\mathcal{F}_{G,m}$.*

By Algorithm B, we immediately obtain the following corollary.

Corollary 2.2 (*Kostic, Yan [12]*) *For any $F \in \mathcal{F}_{G,m}$, let f_F be the (G, m) -multiparking function corresponding with F , i.e., $f_F = \Phi^{-1}(F)$. Let $\sigma(F)$ be the number of the connected components in F and $R(F)$ the set of all the roots in F . Then $\sum_{v \in [n]} f_F(v) = \sum_{e \in E(F)} c(e) + \sum_{v \in V(G) \setminus R(F)} |N(v)| - \sigma(F)$.*

Proof. For any $F \in \mathcal{F}_{G,m}$, let $f_F = \Phi^{-1}(F)$. Recall that $\sigma(F)$ is the number of the connected components in F . For any $v \in [n]$, if v is the minimal vertex which is no less than m in the tree of F , then $f_F(v) = -1$; otherwise, $f_F(v)$ is the sum of the color of the edge e connecting the vertices v to $pre_F(v)$ and the cardinality of the set $N(v)$. Hence,

$$\begin{aligned} \sum_{v \in [n]} f_F(v) &= \sum_{f_F(v) \neq -1} f_F(v) + \sum_{f_F(v) = -1} f_F(v) \\ &= \sum_{f_F(v) \neq -1} f_F(v) - \sigma(F) \\ &= \sum_{e \in E(F)} c_F(e) + \sum_{v \in V(G) \setminus R(F)} |N(v)| - \sigma(F) \end{aligned}$$

■

3 (G, m) -multiparking complement functions

In this section, we will define the (G, m) -multiparking complement function, give the reciprocity theorem for (G, m) -multiparking function. First, let $deg_G(i)$ be the number of edges which are

incident with the vertex i in the graph G . For any subset $I \subseteq V(G)$ and $i \in I$, define $\text{indeg}_I(i)$ as the number of edges from i to vertices inside I . We give the definition of (G, m) -multiparking complement function as follows.

Definition 3.1 Let $m \in [n]$ and G be a connected graph with vertex set $V(G) = [n]$. A (G, m) -multiparking complement function is a function $h : V(G) \rightarrow \mathbb{N}$, such that for every $I \subseteq V(G)$ either $(\overline{A})$ $h(\alpha(I, m)) = \deg_G(\alpha(I, m)) + 1$, or $(\overline{B})$ there exists a vertex $i \in I$ such that $\text{indeg}_I(i) < h(i) \leq \deg_G(i)$.

Given a function $f : V(G) \rightarrow \mathbb{N} \cup \{-1\}$, define a function h_f as $h_f(i) = \deg_G(i) - f(i)$ for all $i \in V(G)$. The following lemma tells us the relation between (G, m) -multiparking functions and (G, m) -multiparking complement functions.

Lemma 3.1 Let $m \in [n]$ and G be a connected graph with vertex set $V(G) = [n]$. Then the function f is a (G, m) -multiparking function if and only if the function h_f is a (G, m) -multiparking complement function.

Proof. Let $f \in \mathcal{MP}_{G,m}$. For any $I \subseteq V(G)$, if I contains no well-behaved vertices, then $f(\alpha(I, m)) = -1$, thus, $h_f(\alpha(I, m)) = \deg_G(\alpha(I, m)) + 1$; otherwise, there exists a vertex $i \in I$ such that $0 \leq f(i) < \text{outdeg}_I(i)$, then $\text{indeg}_I(i) < h_f(i) \leq \deg_G(i)$. Hence, h_f is a (G, m) -multiparking complement function. $\blacksquare$

Let $\overline{\mathcal{MP}}_{G,m}$ be a set of all the (G, m) -multiparking complement functions. The generating function $P_{G,m}(q)$ of (G, m) -multiparking functions is defined as

$$P_{G,m}(q) = \sum_{f \in \mathcal{MP}_{G,m}} q^{\sum_{i \in V(G)} f(i)}$$

Define the generating function of (G, m) -multiparking complement functions to be the polynomial

$$\begin{aligned} \overline{P}_{G,m}(q) &= \sum_{h \in \overline{\mathcal{MP}}_{G,m}} q^{\sum_{i \in V(G)} h(i) - |E(G)|} \\ &= \sum_{f \in \mathcal{MP}_{G,m}} q^{\sum_{i \in V(G)} [\deg_G(i) - f(i)] - |E(G)|} \\ &= q^{|E(G)|} P_{G,m}\left(\frac{1}{q}\right) \end{aligned}$$

Clearly, the sums $\sum_{i \in V(G)} [\deg_G(i) - f(i)] - |E(G)|$ and $\sum_{i \in V(G)} f(i)$ are connected with the reciprocity law for (G, m) -multiparking function. Now, we are in a position to give the reciprocity theorem for (G, m) -multiparking function.

A color forest F on $[n]$ may appear as a subgraph of different graphs, and a vertex function f may be a (G, m) -multiparking function for different graphs. Let $f = \Phi(F)$. Recall that $G - e$ be

a graph obtained by deleting the edge $e = \{i, j\}_k$ from G . We say an edge $e = \{i, j\}_k$ of $G - F$ is *F-redundant* if $\Phi_{G-e, m}(F) = f$. There are the closed relations between (G, m) -multiparking complement functions and *F-redundant* edges of G .

Let π be the order defined in Step 1 of Algorithm *B*. Note that π only depends on F and τ , not the underlying graph G . We have the following lemma.

Lemma 3.2 (*Kostic, Yan [12]*) *Fix a vertex ranking τ . An edge $e = \{v, w\}_k$ of G is F-redundant if and only if e is one of the following types:*

1. *Both v and w are roots of F .*
2. *v is a root and w are non-roots of F , and $\pi^{-1}(w) < \pi^{-1}(v)$.*
3. *v and w are non-roots and $\pi^{-1}(\text{pre}_F(v)) < \pi^{-1}(w) < \pi^{-1}(v)$. In this case v and w must lie in the same tree of F .*
4. *e is a loop of G .*
5. *$e = \{v, \text{pre}_F(v)\}_k$ with $k > k'$, where $\{v, \text{pre}_F(v)\}_{k'} \in F$.*

The results 1, 2 and 3 in Lemma 3.2 were proved in [12]. It is easy to see the results 4 and 5 hold by the Algorithm *A*.

By the above lemma, given a graph G , let F be a spanning color m -subforest of G . Define a function $g_F : V(G) \rightarrow \mathbb{N}$ by letting $g_F(i)$ to be the cardinality of the set $\{e = \{i, j\}_k \mid e \text{ is F-redundant and } \pi^{-1}(j) \leq \pi^{-1}(i)\}$. Define the generating function to be the polynomial

$$I_{G, m}(q) = \sum_{F \in \mathcal{F}_{G, m}} q^{\sum_{i \in V(G)} g_F(i)}$$

Theorem 3.1 (*the reciprocity theorem for (G, m) -multiparking function*)

$$q^{|V(G)|} I_{G, m}(q) = \overline{P}_{G, m}(q) = q^{|E(G)|} P_{G, m}\left(\frac{1}{q}\right).$$

Proof. We only prove the first identity. Note that $g_F(i)$ is the cardinality of the set $\{e = \{i, j\}_k \mid e \text{ is F-redundant and } \pi^{-1}(j) \leq \pi^{-1}(i)\}$. By Corollary 2.2, we have

$$\begin{aligned} |E(G)| &= \sum_{i \in V(G)} g_F(i) + \sum_{f_F(i) \neq -1} |N(i)| + \sum_{e \in F} c_F(e) + |E(F)| \\ &= \sum_{i \in V(G)} g_F(i) + \sum_{f_F(i) \neq -1} |N(i)| + \sum_{e \in F} c_F(e) + |V(G)| - \sigma(F) \\ &= \sum_{i \in V(G)} g_F(i) + \sum_{i \in V(G)} f_F(i) + |V(G)|. \end{aligned}$$

Therefore,

$$\begin{aligned}
\overline{P}_{G,m}(q) &= \sum_{f \in \mathcal{MP}_{G,m}} q^{|E(G)| - \sum_{i \in V(G)} f(i)} \\
&= \sum_{F \in \mathcal{F}_{G,m}} q^{\sum_{i \in V(G)} g_F(i) + V(G)} \\
&= q^{|V(G)|} I_{G,m}(q).
\end{aligned}$$

■

4 The recursion for the generating functions $P_{G,n}(q)$

In this section, we always let $m = n$. In this situation, a spanning color n -subforest of G is a spanning color tree of G with root n since G is connected, and a (G, n) -multiparking function is a G -parking function. Let $\mathcal{T}_G = \{T \mid T \text{ is a spanning color tree of } G\}$ and $\mathcal{P}_G = \{f \mid f \text{ is a } G\text{-parking function}\}$. We write $P_{G,n}(q)$ as $P_G(q)$ for short. There is a bijection $\Phi = \Phi_{G,\tau}$ between the sets $\mathcal{T}_G$ and $\mathcal{P}_G$. For any $T \in \mathcal{T}_G$, define $w_G(T) = \sum_{i=1}^n \Phi(T)(i)$. Then $P_G(q) = \sum_{T \in \mathcal{T}_G} q^{w_G(T)}$.

First, the Algorithms A and B tell us that G -parking functions are independent on the loops in G . So, we obtain the following result.

Lemma 4.1 *Suppose that e is a loop of G . Then $P_G(q) = P_{G-e}(q)$.*

Now, suppose that G has a bridge $e = \{i, j\}_0$. After deleting the edge e , we let G_1 be the subgraph of G such that $\{n, i\} \subseteq V(G_1)$ and G_2 another subgraph of G obtained by letting the label of the vertex j become $n+1$.

Lemma 4.2 *Suppose that $e = \{i, j\}_0$ is a bridge of G . Then $P_G(q) = qP_{G_1}(q)P_{G_2}(q)$.*

Proof. For any $T_1 \in \mathcal{T}_{G_1}$ and $T_2 \in \mathcal{T}_{G_2}$, the trees T_1 and T_2 have the roots n and $n+1$ respectively. Let T be a tree obtained by setting the label of the vertex $n+1$ as j and adding j to be the child of the vertex i . Then T is a spanning color tree of G . Conversely, for any $T \in \mathcal{T}_G$, after deleting the edge e , we let T_1 be the subgraph of T such that $\{n, i\} \subseteq V(T_1)$ and T_2 another subgraph of T obtained by letting the label of the vertex j become $n+1$. Then $T_i \in \mathcal{T}_{G_i}$ for $i = 1, 2$. Take the vertex ranking τ such that $\tau(i) = n-1$ and $\tau(j) = n-2$. Then $w_G(T) = w_{G_1}(T_1) + w_{G_2}(T_2) + 1$ since e is a bridge of G . This implies that

$$\sum_{T \in \mathcal{T}_G} q^{w_G(T)} = q \sum_{T \in \mathcal{T}_{G_1}} q^{w_{G_1}(T)} \sum_{T \in \mathcal{T}_{G_2}} q^{w_{G_2}(T)}.$$

Hence, we have $P_G(q) = qP_{G_1}(q)P_{G_2}(q)$.

■

Now, we consider the case in which $e = \{i, j\}_k$ is neither a loop nor a bridge with $i > j$. Define a graph $G \setminus e$ as follows. The graph $G \setminus e$ is obtained from G contracting the the vertices i and j ; that is, to get $G \setminus e$ we identify two vertices i and j as a new vertex i . First, we discuss the cases in which e is incident with the root n .

Lemma 4.3 *Suppose that $e = \{n, i\}_k \in E(G)$ and e is neither a loop nor a bridge in G . Then $P_G(q) = qP_{G-e}(q) + P_{G \setminus e}(q)$.*

Proof. First, we take the vertex ranking τ such that $\tau(i) = 1$. Let $\mathcal{T}_{G,0} = \{T \in \mathcal{T}_G \mid e \notin T\} \cup \{T \in \mathcal{T}_G \mid e \in T \text{ and } c_T(e) \geq 1\}$ and $\mathcal{T}_{G,1} = \{T \in \mathcal{T}_G \mid e \in T \text{ and } c_T(e) = 0\}$. Given a spanning color tree T of G , let $\pi = \pi_{T,\tau}$ be a permutation $\pi = (\pi(1), \pi(2), \dots, \pi(n)) = (v_1 v_2 \dots v_n)$ on the vertices of G by Step 1 of the Algorithm B. For any $T \in \mathcal{T}_{G,0}$, if $e \notin T$, suppose $j = \text{pre}_T(i)$, then $j \neq n$ and $\pi^{-1}(j) + 1 = \pi^{-1}(i)$ since $\tau(i) = 1$. Thus $w_{G-e}(T) = w_G(T) - 1$ since T is a spanning color tree of $G - e$ as well. If $e \in T$ and $c_T(e) \geq 1$, then let T' be the tree obtained by setting the color of the edge e as $c_T(e) - 1$. The tree T' is a spanning color tree of $G - e$ and $w_{G-e}(T) = w_G(T) - 1$.

For any $T \in \mathcal{T}_{G,1}$, it is easy to see that $\text{pre}_T(i) = n$ and $\pi^{-1}(i) = 2$. Let A be the set of the vertices $j \in \text{child}_T(i)$ such that $n \in N(j)$. Since $c_T(e) = 0$, let T' be the tree obtained by deleting the vertex i , attaching the vertex j to be the child of n and setting the color of the edge $\{n, j\}$ as $c_T(\{i, j\}) + \mu_G(n, j)$ for all $j \in A$. The tree T' can be viewed as a spanning color tree of $G \setminus e$. Clearly, $w_{G \setminus e}(T') = w_G(T)$. Conversely, Given a tree $T' \in \mathcal{T}_{G \setminus e}$, let A be the set of the vertices which is adjacent to n in T such that either $c_T(\{n, j\}) \geq \mu_G(n, j)$ or j isn't adjacent to n in G . Add a new vertex i , attach i to be the child of n and let the color of the edge $\{n, i\}$ be 0. Then delete the edges $\{n, j\}$, attach j to be the child of i and setting the color of $\{n, j\}$ as $c_T(\{n, j\}) - \mu_G(n, j)$ for all $j \in A$. The obtained tree can be viewed as a spanning color tree of G . Hence, $P_G(q) = qP_{G-e}(q) + P_{G \setminus e}(q)$. $\blacksquare$

Next, we discuss the case in which e isn't incident with the root n . Suppose $e = \{i, j\}_k \in E(G)$ with $i > j$. Take the vertex ranking τ such that $\tau(i) = 1$ and $\tau(j) = 2$. Define the following six sets:

- (1) $\hat{T}_{G,0}^1 = \{T \in \mathcal{T}_G \mid e \notin T \text{ and } \text{pre}_T(j) \neq \text{pre}_T(i)\}$
- (2) $\hat{T}_{G,0}^2 = \{T \in \mathcal{T}_G \mid e \notin T, \text{pre}_T(j) = \text{pre}_T(i) \text{ and } c_T(\{j, \text{pre}_T(i)\}) \neq 0\}$
- (3) $\hat{T}_{G,0}^3 = \{T \in \mathcal{T}_G \mid e \notin T, \text{pre}_T(j) = \text{pre}_T(i) \text{ and } c_T(\{j, \text{pre}_T(i)\}) = 0\}$
- (4) $\hat{T}_{G,1}^1 = \{T \in \mathcal{T}_G \mid e \in T \text{ and } c_T(e) \geq 1\}$
- (5) $\hat{T}_{G,1}^2 = \{T \in \mathcal{T}_G \mid e \in T, \pi_T^{-1}(i) < \pi_T^{-1}(j), \text{pre}_T(i) \in N_{G,T}(j) \text{ and } c_T(e) = 0\}$, where $N_{G,T}(v) = \{v_j \mid \{v, v_j\}_k \in E(G) \text{ and } \pi^{-1}(v_j) < \pi^{-1}(\text{pre}_F(v))\}$.
- (6) $\hat{T}_{G,1}^3 = \{T \in \mathcal{T}_G \mid e \in T\} \setminus (\hat{T}_{G,1}^1 \cup \hat{T}_{G,1}^2)$.

Lemma 4.4 *Suppose that $e = \{i, j\}_k \in E(G)$ and e is neither a loop nor a bridge in G . There is a bijection ψ from $\hat{T}_{G,0}^1 \cup \hat{T}_{G,0}^2 \cup \hat{T}_{G,1}^1 \cup \hat{T}_{G,1}^2$ to $\mathcal{T}_{G-e}$. Moreover, $w_{G-e}(\psi(T)) = w_G(T) - 1$ for any $T \in \hat{T}_{G,0}^1 \cup \hat{T}_{G,0}^2 \cup \hat{T}_{G,1}^1 \cup \hat{T}_{G,1}^2$.*

Proof. For any $T \in \hat{T}_{G,0}^1$, let $\psi(T) = T$. $\psi(T)$ can be viewed as a spanning tree of $G - e$. Moreover, $w_{G-e}(T) = w_G(T) - 1$.

For any $T \in \hat{T}_{G,0}^2$, suppose $v = \text{pre}_T(i)$. Since $c_T(\{j, v\}) \neq 0$, let T' be the tree obtained by setting the color of $\{j, v\}$ as $c_T(\{j, v\}) - 1$. The tree T' can be viewed as a spanning tree of $G - e$ and satisfies that $\text{pre}_{T'}(i) = \text{pre}_{T'}(j)$ and $c_{T'}(\{j, v\}) \leq \mu_{G-e}(j, v) - 2 = \mu_G(j, v) - 2$. Moreover, $w_{G-e}(T') = w_G(T) - 1$.

For any $T \in \hat{T}_{G,1}^1$, let T' be the tree obtained by setting the color of $\{i, j\}$ as $c_T(i, j) - 1$. The tree T' can be viewed as a spanning tree of $G - e$ and $c_{T'}(\{i, j\}) \geq 0$. Moreover, $w_{G-e}(T') = w_G(T) - 1$.

For any $T \in \hat{T}_{G,1}^2$, suppose that $v = \text{pre}_T(i)$. Let T' be the tree obtained by deleting the edge $\{i, j\}_0$, attaching j to be the child of v and setting the color of the edge $\{v, j\}$ as $\mu_G(v, j) - 1$. Since $v \in N(j)$, the tree T' can be viewed as a spanning tree of $G - e$, $c_{T'}(\{j, v\}) = \mu_{G-e}(j, v) - 1 = \mu_G(j, v) - 1$ and $\text{pre}_{T'}(i) = \text{pre}_{T'}(j)$. Moreover, $w_{G-e}(T') = w_G(T) - 1$. This complete the proof. ■

Lemma 4.5 Suppose that $e = \{i, j\}_k \in E(G)$ and e is neither a loop nor a bridge in G . There is a bijection ψ' from $\hat{T}_{G,0}^3 \cup \hat{T}_{G,1}^3$ to $\mathcal{T}_{G \setminus e}$. Moreover, $w_{G \setminus e}(\psi(T)) = w_G(T)$ for any $\hat{T}_{G,0}^3 \cup \hat{T}_{G,1}^3$.

Proof. For any $T \in \hat{T}_{G,0}^3$, suppose $v = \text{pre}_T(j)$. Let T' be the tree obtained by deleting the vertex j , attaching the vertex w to be the child of i and setting the color of $\{i, w\}$ as $c_T(\{j, w\}) + \mu_G(i, w)$ for any $w \in \text{child}_T(j)$. The tree T' can be viewed as a spanning tree of $G \setminus e$ and $0 \leq c_{T'}(\{i, v\}) \leq \mu_G(\{i, v\}) - 1$. Moreover, $w_{G \setminus e}(T') = w_G(T)$.

For any $T \in \hat{T}_{G,1}^3$, we always have $c_T(\{i, j\}) = 0$. we discuss the following three cases:

Case 1. $\pi_T^{-1}(i) < \pi_T^{-1}(j)$ and $\text{pre}_T(i) \notin N(j)$

Let T' be the tree obtained by deleting the vertex j , attaching w to be the child of i and setting the color of the edge $\{w, i\}$ as $c_T(\{j, w\}) + \mu_G(i, w)$ for all $w \in \text{child}_T(j)$. The tree T' can be viewed as a spanning tree of $G \setminus e$. Moreover, $w_{G \setminus e}(T') = w_G(T)$.

Case 2. $\pi_T^{-1}(i) > \pi_T^{-1}(j)$ and $\text{pre}_T(j) \notin N(i)$

Suppose $v = \text{pre}_T(j)$. Let T' be the tree obtained by deleting the vertex j , attaching i to be the child of v , attaching w to be the children of i for all $w \in \text{child}_T(j)$ and setting the color of the edge $\{u, i\}$ as $c_T(\{i, u\}) + \mu_G(j, u)$ for all $u \in \text{child}_T(i)$. The tree T' can be viewed as a spanning tree of $G \setminus e$. Moreover, $w_{G \setminus e}(T') = w_G(T)$.

Case 3. $\pi_T^{-1}(i) > \pi_T^{-1}(j)$ and $\text{pre}_T(j) \in N_{G,T}(i)$

Suppose that $v = \text{pre}_T(j)$. Let T' be the tree obtained by deleting the vertex j , attaching i to be the child of v , attaching w to be the children of i for all $w \in \text{child}_T(j)$ and setting the color of the edge $\{w, i\}$ as $c_T(\{i, w\}) + \mu_G(j, w)$ for all $w \in \text{child}_T(i)$ and the color of the edge $\{v, i\}$ as $c_T(\{j, v\}) + \mu_G(i, v)$. The tree T' can be viewed as a spanning tree of $G \setminus e$ and $c_{T'}(\{i, v\}) \geq \mu_G(i, v)$. Moreover, $w_{G-e}(T') = w_G(T)$.

Conversely, for any $T \in \mathcal{T}_{G \setminus e}$, suppose $v = \text{pre}_T(i)$. we consider the following three cases.

Case 1. $\mu_G(i, v) \geq 1$, $\mu_G(j, v) \geq 1$ and $0 \leq c_T(\{i, v\}) \leq \mu_G(i, v) - 1$.

Add a new vertex j and attach j to be the child of v . Let the color of $\{v, j\}$ be 0. For any $w \in \text{child}_T(i)$, if $c_T(\{i, w\}) \geq \mu_G(i, w)$ or $\mu_G(i, w) = 0$, then delete the edge $\{i, w\}$ and attach w to be the child of j , let the color of $\{j, w\}$ be $c_T(\{i, w\}) - \mu_G(i, w)$. The obtained tree can be view as a spanning tree of G .

Case 2. $\mu_G(i, v) \geq 1$, $\mu_G(j, v) \geq 1$ and $c_T(\{i, v\}) \geq \mu_G(i, v)$

Add a new vertex j and attach j to be the child of v . Delete the edge $\{v, i\}$ and attach i to be the child of j . Let the color of $\{v, j\}$ be $c_T(\{i, v\}) - \mu_G(i, v)$ and the color of $\{i, j\}$ 0. For any $w \in \text{child}_T(i)$, if $c_T(\{i, w\}) \geq \mu_G(j, w)$ or $\mu_G(j, w) = 0$, then let the color of the edges $\{i, w\}$ be $c_T(\{i, w\}) - \mu_G(j, w)$; otherwise delete the edge $\{i, w\}$ and attach w to be the child of j . The obtained tree can be view as a spanning tree of G .

Case 3. There is exact one vertex w such that $\mu_G(w, v) \geq 1$ for $w \in \{i, j\}$

We first consider the case $\mu_G(i, v) \geq 1$, $\mu_G(j, v) = 0$. Add a new vertex j and attach j to be the child of i . Let the color of $\{i, j\}$ be 0. For any $w \in \text{child}_T(i)$, if $c_T(\{i, w\}) \geq \mu_G(i, w)$ or $\mu_G(i, w) = 0$, then delete the edge $\{i, w\}$ and attach w to be the son of j . Let the color of $\{j, w\}$ be $c_T(\{i, w\}) - \mu_G(i, w)$. The obtained tree can be view as a spanning tree of G . Similarly, we may consider the case $\mu_G(i, v) = 0$, $\mu_G(j, v) \geq 1$. The proof is completed. ■

Theorem 4.1 *Suppose that $e \in E(G)$ and e is neither a loop nor a bridge in G . Then*

$$P_G(q) = qP_{G-e}(q) + P_{G \setminus e}(q).$$

Proof. Combining Lemmas 4.4 and 4.5, we obtain the results as desired. ■

References

- [1] R. Cori, D. Rossin, B. Salvy, Polynomial ideals for sandpiles and their Grobner bases. *Theoretical Computer Science* **276** (2002), no. 1-2, 1-15.
- [2] Chebikin Denis, Pylyavskyy Pavlo, A family of bijections between G -parking functions and spanning trees. *J. Combin. Theory Ser. A* **110** (2005), no. 1, 31–41.
- [3] Dhar, D. Self-organized critical state of the sandpile automaton models, *Physical Review Letters* 64(1990), no. 14, 1613-1616.
- [4] Sen-Peng Eu, Tung-Shan Fu, Chun-Ju Lai, On the enumeration of parking functions by leading terms, *Adv. in Appl. Math.* 35 (2005) 392-406
- [5] Sen-Peng Eu, Tung-Shan Fu, Yeong-Nan Yeh, Refined Chung-Feller theorems for lattice paths, *J. Combin. Theory, Ser. A* 112 (2005) 143-162.
- [6] Sen-Peng Eu, Shu-Chung Liu, Yeong-Nan Yeh, Taylor expansions for Catalan and Motzkin numbers, *Adv. in Appl. Math.* 29 (2002) 345-357.

- [7] D. Foata, J. Riordan, Mappings of acyclic and parking functions, *Aequationes Math.* 10 (1974) 10-22.
- [8] J. Françon, Acyclic and parking functions, *J. Combin. Theory Ser. A* 18 (1975) 27-35.
- [9] J.D. Gilbey, L.H. Kalikow, Parking functions, valet functions and priority queues, *Discrete Math.* 197/198 (1999) 351-373.
- [10] D. Knuth, Linear probing and graphs, average-case analysis for algorithms, *Algorithmica* 22 (4) (1998) 561C568.
- [11] Konheim, A. G. and Weiss, B. An Occupancy Discipline and Applications. *Siam Journal of Applied Mathematics* 14 (1966) 1266-1274
- [12] Kostić Dimitrije, Yan, Catherine H. Multiparking functions, graph searching, and the Tutte polynomial. *Adv. in Appl. Math.* 40 (2008), no. 1, 73–97.
- [13] G. Kreweras, Une famille de polynômes ayant plusieurs propriétés énumératives, *Period. Math. Hungar.* 11 (1980) 309-320.
- [14] J.P.S. Kung, C.H. Yan, Gončarov polynomials and parking functions, *J. Combin. Theory Ser. A* 102 (2003) 16-37.
- [15] Postnikov, A. and Shapiro, B. Trees, Parking Functions, Syzygies, and Deformations of Monomial Ideals. *Transactions of the American Mathematical Society* 356 (2004).
- [16] J. Pitman, R. Stanley, A polytope related to empirical distributions, plane trees, parking functions, and the associahedron, *Discrete Comput. Geom.* 27 (4) (2002) 603-634.
- [17] J.Riordan, Ballots and trees, *J. Combin. Theory* 6 (1969) 408-411.
- [18] M.P. Schützenberger, On an enumeration problem, *J. Combin. Theory* 4 (1968) 219-221.
- [19] R.P. Stanley, Hyperplane arrangements, interval orders and trees, *Proc. Natl. Acad. Sci.* 93 (1996) 2620-2625.
- [20] R.P. Stanley, Parking functions and non-crossing partitions, in: The Wilf Festschrift, *Electron. J. Combin.* 4 (1997) R20.
- [21] R.P. Stanley, Hyperplane arrangements, parking functions, and tree inversions, in: B. Sagan, R. Stanley (Eds.), *Mathematical Essays in Honor of Gian-Carlo Rota*, Birkhäuser, Basel, 1998, pp. 359-375.
- [22] R. Stanley, *Enumerative Combinatorics, Volume I*, Cambridge University Press, 1997.

- [23] C.H. Yan, Generalized tree inversions and k-parking functions, *J. Combin. Theory Ser. A* 79 (1997) 268-280.
- [24] C.H. Yan, On the enumeration of generalized parking functions, *Congr. Numer.* 147 (2000) 201-209.
- [25] C.H. Yan, Generalized parking functions, tree inversions and multicolored graphs, *Adv. in Appl. Math.* 27 (2001) 641-670.